arXiv:1010.3871v1 [math.RT] 19 Oct 2010

QUIVERS WITHOUT LOOPS ADMIT GLOBAL DIMENSION 2

NICOLAS POETTERING

Abstract. Let Q be a finite quiver without loops. Then there is an admissible ideal I such that the algebra kQ/I has global dimension at most two and is (strongly) quasi-hereditary. In addition some other (strongly) quasi-hereditary algebras kQ/I' are constructed with bigger global dimension.

1. Introduction

Let Q be a finite quiver and I an admissible ideal of the path algebra kQ. The global dimension $\operatorname{gldim}(kQ/I)$ of the finite-dimensional algebra kQ/I is an important invariant of the category of finite-dimensional left kQ/I-modules. We are interested in the following question: Which natural numbers can actually occur as global dimensions of an algebra kQ/I for some fixed finite quiver Q with different admissible ideals I?

It is well known that if Q contains a loop, the global dimension of kQ/I is infinite for any admissible ideal I. For a finite quiver Q with at least one arrow an admissible ideal I with $\operatorname{gldim}(kQ/I) = 1$ exists if and only if Q has no oriented cycles. In Corollary 2 we answer this question for global dimension 2. But we only consider small global dimensions since the global dimension of a finite-dimensional algebra is neither bounded by a function depending on the number of simple modules (see [1]) nor by a function on the Loewey length of the algebra (see [2]).

1.1. Main Result.

A quiver $R = (R_0, R_1, s_R, t_R)$ is a *subquiver* of a quiver $Q = (Q_0, Q_1, s_Q, t_Q)$, if $R_0 \subseteq Q_0$, $R_1 \subseteq Q_1$, $s_R = s_Q|_{R_1}$ and $t_R = t_Q|_{R_1}$. For $m \in \mathbb{N}_{>0}$ let A_m be the quiver

$$\begin{tikzcd} 1 \arrow[r] & 2 \arrow[r] & \cdots \arrow[r] & m-1 \arrow[r] & m \end{tikzcd}$$

and X_m the quiver

$$\begin{tikzcd} 1 \arrow[r] & 2 \arrow[r] & \cdots \arrow[r] & i \arrow[r] & \cdots \arrow[r] & m \arrow[ll, bend right] \end{tikzcd}$$

with some $1 \leq i < m$.

Theorem 1. *Let Q be a finite quiver without loops.*

(*a*) *Then there exists an admissible ideal I such that* $\operatorname{gldim}(kQ/I) \leq 2$.

(*b*) *Let $k, m \in \mathbb{N}$ with $2 \leq k < m$. If A_m is a subquiver of Q, then there exists an admissible ideal I' such that* $\operatorname{gldim}(kQ/I') = k$.

(*c*) *Let $k, m \in \mathbb{N}$ with $2 \leq k \leq m$. If X_m is a subquiver of Q, then there exists an admissible ideal I' such that* $\operatorname{gldim}(kQ/I') = k$.

Moreover all these ideals are generated by zero relations of length at most three and the algebras are strongly quasi-hereditary.

The proof of this theorem is given in Section 3. The following corollary is a direct consequence.

Corollary 2. *Let Q be a finite quiver. Then there is an admissible ideal I such that* $\operatorname{gldim}(kQ/I) = 2$ *if and only if Q has no loops and* $(kQ^+)^2 \neq 0$.

Supported by BIGS-Mathematics, Bonn and Mathematical Institute of the University Bonn.

Example 3. Let Q be the quiver



I the admissible ideal $\langle da, cf, ef, cb, eb\rangle$ and $I' = I + \langle ba\rangle$. Then $\mathrm{gldim}(kQ/I) = 2$, $\mathrm{gldim}(kQ/I') = 3$ and the algebras kQ/I and kQ/I' are strongly quasi-hereditary.

1.2. **Acknowledgement.** I thank Dieter Happel for asking this nice question. He obtained most of these results independently. Additionally I thank Martin Kalck for discussing and reading my results.

2. Definitions

2.1. **Path algebra.** Let k be a field. Let $Q = (Q_0, Q_1, s, t)$ be a *finite quiver*, i.e. a finite oriented graph with vertex set Q_0, arrow set Q_1 and maps $s, t\colon Q_1 \to Q_0$ indicating the start and terminal point of each arrow. An *oriented path* $\rho = a_1 \dots a_n$ in Q is the concatenation of some arrows $a_1, \dots, a_n \in Q_1$ such that $t(a_{i+1}) = s(a_i)$ for all $1 \leq i < n$. Additionally we introduce a path e_i of length zero for each vertex $i \in Q_0$. The *path algebra* kQ of a quiver Q is the k-vector space with the set of oriented paths as a basis. The product of basis vectors is given by the concatenation of paths if possible or by zero otherwise. Let kQ^+ be the ideal in the path algebra kQ, which is generated by all arrows in Q_1. An two sided ideal I of the path algebra kQ is called *admissible* if there is a $k \in \mathbb{N}_{>0}$ with $(kQ^+)^k \subseteq I \subseteq (kQ^+)^2$.

Let Q be a finite quiver and I an admissible ideal. Then kQ/I is a finite-dimensional k-algebra. The isomorphism classes of simple left kQ/I-modules are in a unique bijection with the vertices Q_0 of the quiver Q. So we denote the simple module associated to $i \in Q_0$ by $S(i)$ and the projective cover of $S(i)$ by $P(i)$.

2.2. **Global dimension.** Let Q be a finite quiver, I an admissible ideal and M a kQ/I-module. The *projective dimension* $\mathrm{pdim}(M)$ of the module M is the length of a minimal projective resolution. The *global dimension* $\mathrm{gldim}(kQ/I)$ of the algebra kQ/I is the supremum of the projective dimensions of all modules. These dimensions are in general not finite. But it is well known that the global dimension of kQ/I is the maximum of the projective dimensions of all simple modules.

2.3. **Quiver of a module.** Let Q be a finite quiver, I an admissible ideal and M a kQ/I-module. Take some Jordan Hölder filtration $0 = M_0 \subseteq M_1 \subseteq M_2 \subseteq \dots \subseteq M_t = M$ of M with $M_i/M_{i-1} \cong S(j(i))$ and $j(i) \in Q_0$ for all $1 \leq i \leq t$. Thus for each $1 \leq i \leq t$ a vector $v_i \in M_i - M_{i-1}$ exists such that $e_{j(i)}v_i = v_i$. Now we associate a quiver to the module M:

The vertices are the basis vectors $v_1, \dots, v_t$. Let v_i and v_j be two vectors and $a \in Q_1$ such that $\lambda_j \neq 0$ in $av_i = \sum_k \lambda_k v_k$. In this case there is one arrow called a from v_i to v_j. Note that in general this quiver depends on the chosen basis.

For the quiver associated to M we write $j(i)$ instead of v_i for all $1 \leq i \leq t$.

2.4. **Quasi-hereditary algebra.** The algebras constructed in the proof of Theorem 1 are strongly quasi-hereditary in the sense of Ringel [3].

Definition 4. Let Q be a finite quiver with $Q_0 = \{1, 2, \dots, n\}$ and I an admissible ideal. We say that kQ/I is *strongly quasi-hereditary* if there is for any $i \in Q_0$ an exact sequence

$$0 \to R(i) \to P(i) \to \Delta(i) \to 0 \tag{1}$$

with the following two properties:

(a) $R(i)$ is a direct sum of projective modules $P(j)$ with $j < i$.
(b) If $S(j)$ is a composition factor of $\operatorname{rad}(\Delta(i))$, then $j > i$.

In [3] Ringel proved that a strongly quasi-hereditary algebra kQ/I with $Q_0 = \{1, 2, \dots, n\}$ is also quasi-hereditary and has global dimension at most n.

2.5. Examples.

Example 5. Let Q be the quiver with one vertex 1 and one loop a and let $I = \langle a^3 \rangle$. Then the quiver associated to $S(1)$ and $P(1)$ are given by

$$1 \qquad 1 \xrightarrow{a} 1 \xrightarrow{a} 1.$$

In this case $\operatorname{pdim} S(1) = \infty$, $\operatorname{pdim} P(1) = 0$ and $\operatorname{gldim}(kQ/I) = \infty$. Thus the algebra kQ/I is not strongly quasi-hereditary.

Example 6. Let $n \in \mathbb{N}_{>0}$, Q the quiver $1 \xrightarrow{a_1} 2 \xrightarrow{a_2} \dots \xrightarrow{a_{n-1}} n$ and $I = \langle a_{i+1}a_i | 1 \leq i \leq n-2 \rangle$. Then the quivers of $S(i)$ and $P(j)$ with $i, j \in Q_0$ and $j \neq n$ are given by

$$i \qquad j \xrightarrow{a_j} j+1.$$

Then $\operatorname{pdim} S(i) = n - i$, $\operatorname{gldim}(kQ/I) = n-1$ and kQ/I is strongly quasi-hereditary.

3. Proof of Theorem 1

The proof of the main theorem is divided into five propositions. The following relabeling of the vertices and arrows of Q is possible for any finite quiver without loops. Let $n, r_{ij} \in \mathbb{N}$ and Q be the quiver with

$$\begin{aligned} Q_0 &= \{1, \dots, n\}, \\ Q_1 &= \{a_{ijt} \colon i \to j | i, j \in Q_0, t \in \mathbb{N}, i \neq j, 1 \leq t \leq r_{ij}\}. \end{aligned}$$

Let I be the ideal

$$I = \langle a_{kit} a_{jku} | i, j, k \in Q_0, t, u \in \mathbb{N}, i < k, j < k, 1 \leq t \leq r_{ki}, 1 \leq u \leq r_{jk} \rangle. \tag{2}$$

This ideal is admissible, since $(kQ^+)^{2n-1} \subseteq I$.

Proposition 7. *Let Q be a finite quiver without loops and I the ideal defined in Equation* (2)*. Then* $\operatorname{gldim}(kQ/I) \leq 2$.



Figure 1. The projective modules $P(i)$ with $i \in \{1, 2, 3, 4\}$ for $n = 4$ and $r_{ij} = 1$.

Proof. Figure 1 shows the quivers of the indecomposable projective modules for $n = 4$ and $r_{ij} = 1$ for all $i, j \in Q_0$ with $i \neq j$. Using these pictures we get in general for each $i \in Q_0$ the following minimal projective resolution:

$$0 \to \bigoplus_{j \in Q_0, j > i} \bigoplus_{k \in Q_0, k < j} P(k)^{r_{ij} r_{jk}} \to \bigoplus_{j \in Q_0 - \{i\}} P(j)^{r_{ij}} \to P(i) \to S(i) \to 0. \tag{3}$$

This means $\operatorname{gldim}(kQ/I) \leq 2$. □

Let A_m be a subquiver of Q. Then A_k with $k \leq m$ is also a subquiver of Q. If A_m can be extended to some X_m in Q (i.e. there is a $a \in Q_1$ with $s(a) = m$ and $t(a) \in (A_m)_0$), then it is enough to study the case that X_m is a subquiver of Q (see Proposition 11). Otherwise Part (b) of Theorem 1 is proven in the following proposition.

Proposition 8. *Let Q be a finite quiver without loops and $m \in \mathbb{N}$ with $m \geq 3$. If A_m is a subquiver of Q, which cannot be extended to some X_m in Q, then there exists an admissible ideal I' such that* $\operatorname{gldim}(kQ/I') = m - 1$.

Proof. Let $m \in \mathbb{N}$ with $m \geq 3$ and A_m a subquiver of Q. Thus we have $r_{i,i+1} > 0$ for all $1 \leq i < m$. We use the ideal I defined in (2) and Proposition 7. Let

$$\begin{aligned} I' = I + \langle a_{i+1,i+2,t}a_{i,i+1,u} | i \in Q_0, t, u \in \mathbb{N}, i \leq m-2, \\ 1 \leq t \leq r_{i+1,i+2}, 1 \leq u \leq r_{i,i+1} \rangle. \end{aligned}$$

For $i \in Q_0$ define $R(i) = \bigoplus_{j \in Q_0, j<i} P(j)^{r_{ij}}$ and $\Delta(i) = P(i)/R(i)$ using that $R(i)$ is a direct summand of $\operatorname{rad} P(i)$. For $i \in Q_0$ with $i < m$ let $\Gamma(i)$ be the cokernel of some map $R(i) \oplus P(i+1)^{r_{i,i+1}} \to P(i)$ such that $\Gamma(i)$ is minimal with this property. Thus $\Gamma(i)$ is unique and the top of $\Gamma(i)$ is isomorphic to $S(i)$.

Again we give the quivers of indecomposable projective modules for $n = 5$, $m = 4$ and $r_{ij} = 1$ for all $i, j \in Q_0$ with $i \neq j$ in Figure 2. Figure 3 and 4 shows the quivers of the modules $\Delta(i)$ and the $\Gamma(i)$ in this case.



Figure 2. The projective modules $P(i)$ with $i \in \{1, 2, 3, 4, 5\}$ for $n = 5$, $m = 4$ and $r_{ij} = 1$.



Figure 3. The modules $\Delta(i)$ with $i \in \{1, 2, 3, 4, 5\}$ for $n = 5$, $m = 4$ and $r_{ij} = 1$.

Since Sequence (3) is exact for all $i \in Q_0$ with $i \geq m - 1$ we get $\operatorname{pdim}(S(i)) \in \{0, 1, 2\}$ in this case. With Figure 2, 3 and 4 we get in general for each $i \in Q_0$ with $i \leq m - 2$ an exact sequence

$$0 \to R(i) \oplus \Gamma(i+1)^{r_{i,i+1}} \oplus \bigoplus_{j \in Q_0, j \geq i+2} \Delta(j)^{r_{ij}} \to P(i) \to S(i) \to 0. \tag{4}$$



FIGURE 4. The modules $\Gamma(i)$ with $i \in \{1, 2, 3\}$ for $n = 5$, $m = 4$ and $r_{ij} = 1$.

Using $\operatorname{pdim}(\Gamma(i+1)) \geq 1$ for all $i \in Q_0$ with $i \leq m-2$ we get $\operatorname{pdim}(S(i)) = \operatorname{pdim}(\Gamma(i+1)) + 1$. Since in this case the sequence

$$0 \to R(i) \oplus \Gamma(i+1)^{r_{i,i+1}} \to P(i) \to \Gamma(i) \to 0 \tag{5}$$

is exact we know $\operatorname{pdim}(\Gamma(i)) = \operatorname{pdim}(\Gamma(i+1)) + 1$. Thus for $i \in Q_0$ with $i \leq m-2$

$$\operatorname{pdim}(S(i)) = \operatorname{pdim}(\Gamma(i+1)) + 1 = \operatorname{pdim}(\Gamma(m-1)) + m - 1 - i.$$

Therefore $\operatorname{pdim}(S(i)) \leq \operatorname{pdim}(S(1))$ for all $i \in Q_0$ and

$$\operatorname{gldim}(kQ/I') = \operatorname{pdim}(\Gamma(m-1)) + m - 2.$$

It is not hard to compute the minimal projective resolution of $\Gamma(m-1)$:

$$0 \to R(m)^{r_{m-1,m}} \to R(m-1) \oplus P(m)^{r_{m-1,m}} \to P(m-1) \to \Gamma(m-1) \to 0. \tag{6}$$

Now we assume that A_m cannot be extended to some X_m in Q. Thus $R(m) = 0$. Then Sequence (6) provides $\operatorname{pdim}(\Gamma(m-1)) = 1$ and $\operatorname{gldim}(kQ/I') = m - 1$. ☐

Remark 9. The proof of Proposition 8 holds up to Sequence (6) for general subquivers A_m. Hence we can use it for X_m, too.

Let X_m be a subquiver of Q and $i \in (X_m)_0$ with $r_{mi} \neq 0$. So we can assume that i is chosen maximal for fixed vertices $(X_m)_0 = \{1, \ldots, m\}$. Thus (by relabeling the vertices) X_{m-j} is also a subquiver of Q for all $0 \leq j < i$ and A_{m-i} as well. Using Proposition 8 the following two propositions yield Part (c) of Theorem 1.

Proposition 10. *Let Q be a finite quiver without loops and $m \in \mathbb{N}$ with $m \geq 2$. If X_m is a subquiver of Q, then there exists an admissible ideal I' such that $\operatorname{gldim}(kQ/I') = m$.*

Proof. Using the proof of Proposition 8, especially the notation, $R(m) \neq 0$. Then in this case Sequence (6) yields $\operatorname{pdim}(\Gamma(m-1)) = 2$ and $\operatorname{gldim}(kQ/I') = m$. ☐

Proposition 11. *Let Q be a finite quiver without loops and $m \in \mathbb{N}$ with $m \geq 4$. If X_m is a subquiver of Q, then there exists an admissible ideal I'' such that $\operatorname{gldim}(kQ/I'') = m - 1$.*

Proof. This proof is done very similar to the proof of Proposition 8. Let

$$\begin{aligned} I'' = I + \langle a_{i+1,i+2,t}a_{i,i+1,u} | i \in Q_0, t, u \in \mathbb{N}, i \leq m-4, \\ 1 \leq t \leq r_{i+1,i+2}, 1 \leq u \leq r_{i,i+1} \rangle \\ + \langle a_{m-1,m,s}a_{m-2,m-1,t}a_{m-3,m-2,u} | s, t, u \in \mathbb{N}, 1 \leq s \leq r_{m-1,m}, \\ 1 \leq t \leq r_{m-2,m-1}, 1 \leq u \leq r_{m-3,m-2} \rangle. \end{aligned}$$

We define $R(i)$, $\Delta(i)$ and $\Gamma(i)$ as in the proof of Proposition 8. Thus $r_{i,i+1} > 0$ for all $1 \leq i < m$ and $R(m) \neq 0$. Again we give some quivers of indecomposable projective modules for $n = 5$, $m = 4$ and $r_{ij} = 1$ for all $i, j \in Q_0$ with $i \neq j$ in Figure 5.



FIGURE 5. The projective modules $P(1)$ and $P(2)$ for $n = 5$, $m = 4$ and $r_{ij} = 1$.

Since Sequence (3) is exact for all $i \in Q_0$ with $i \geq m - 2$ we get $\operatorname{pdim}(S(i)) \in \{0, 1, 2\}$ in this case. For $i = m - 3$ we construct the following exact sequences with $r = r_{m-3,m-2} r_{m-2,m-1} r_{m-1,m}$:

$$\begin{aligned} 0 &\to R(m)^r \to R(m-2)^{r_{m-3,m-2}} \oplus P(m)^r \\ &\to R(m-3) \oplus P(m-2)^{r_{m-3,m-2}} \oplus \bigoplus_{j \in Q_0, j \geq m-1} \Delta(j)^{r_{m-3,j}} \\ &\to P(m-3) \to S(m-3) \to 0, \\ 0 &\to R(m)^r \to R(m-2)^{r_{m-3,m-2}} \oplus P(m)^r \\ &\to R(m-3) \oplus P(m-2)^{r_{m-3,m-2}} \to P(m-3) \to \Gamma(m-3) \to 0. \end{aligned}$$

Since $\operatorname{pdim}(\Delta(j)) \leq 1$ and $R(m) \neq 0$ we get $\operatorname{pdim}(S(m-3)) = \operatorname{pdim}(\Gamma(m-3)) = 3$. With Figure 5 Sequence (4) and (5) are exact again for each $i \in Q_0$ with $i \leq m-4$. Thus in this case $\operatorname{pdim}(\Gamma(i)) = \operatorname{pdim}(\Gamma(i+1)) + 1$ and

$$\operatorname{pdim}(S(i)) = \operatorname{pdim}(\Gamma(i+1)) + 1 = \operatorname{pdim}(\Gamma(m-3)) + m - 3 - i = m - i.$$

Therefore $\operatorname{gldim}(kQ/I'') = m - 1$. ☐

The following proposition completes the proof of Theorem 1.

Proposition 12. *The algebras occurring in Proposition 7, 8, 10 and 11 are strongly quasi-hereditary.*

Proof. Define $R(i)$ and $\Delta(i)$ as above. Thus Sequence (1) is exact and by the construction of the ideal I holds $\dim_k \operatorname{Hom}_{kQ/I}(P(j), \Delta(i)) = \delta_{ij}$ for all $j \in Q_0$ with $j \leq i$. ☐

REFERENCES

[1] E.L. Green, *Remarks on projective resolutions*, Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), pp. 259-279, Lecture Notes in Math., 832, Springer, Berlin, 1980.

[2] E. Kirkman and J. Kuzmanovich, *Algebras with large homological dimensions*, Proc. Amer. Math. Soc. 109 (1990), no. 4, 903-906.

[3] C.M. Ringel, *Iyama's finiteness theorem via strongly quasi-hereditary algebras* J. Pure Appl. Algebra 214 (2010), no. 9, 1687-1692.

NICOLAS POETTERING, MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, D-53115 BONN, GERMANY

E-mail address: `n.poettering@gmail.com`